

What is the nature of RX J0720.4-3125?

Jeremy S. Heyl
jshey1@ucolick.org

Lars Hernquist¹
lars@ucolick.org

Lick Observatory, University of California, Santa Cruz, California 95064, USA

ABSTRACT

RX J0720.4-3125 has recently been identified as a pulsating soft X-ray source in the ROSAT all-sky survey with a period of 8.391 s. Its spectrum is well characterized by a black-body with a temperature of 8×10^5 K. We propose that the radiation from this object is thermal emission from a cooling neutron star. For this black-body temperature we can obtain a robust estimate of the object's age of $\sim 3 \times 10^5$ yr, yielding a polar field $\sim 10^{14}$ G for magnetic-dipole spin down and a value of $\dot{P}$ compatible with current observations.

Subject headings: stars: neutron — stars: magnetic fields — radiative transfer — X-rays: stars

1. Introduction

“Breaking” (Mereghetti & Stella 1995) or “anomalous” (Van Paradijs, Taam & van den Heuvel 1995) x-ray pulsars (AXPs) typically have pulsed X-ray emission with steadily increasing periods ~ 10 s, X-ray luminosities $\sim 10^{35} - 10^{36}$ erg/s, soft spectra, and no detected companions or accretion disks. Haberl et al. (1997) have recently identified RX J0720.4-3125 as a pulsating, soft X-ray source with a period of 8.391 s. They estimate its bolometric luminosity to be $2.6 \times 10^{31} d_{100}^2$ erg/s, where d_{100} is the object's distance in units of 100 pc.

Wang (1997) has proposed that accretion onto a weakly magnetized neutron star powers RX J0720.4-3125, and that for the neutron star to spin down to 8.391 s within a Hubble time, it either must have been born with a period ~ 0.5 s or have experienced magnetic field decay. Our proposal begins with the second footnote of Wang (1997) which suggests the possibility that RX J0720.4-3125 is powered by an internal heat source. We propose that the neutron-star cooling powers RX J0720.4-3125. This would give it an age $\sim 3 \times 10^5$ yr, much younger than the Wang's (1997) estimated age $\gtrsim 10^9$ yr.

2. Calculations

In a previous paper (Heyl & Hernquist 1997c), we have argued that AXPs may be powered by neutron star cooling through an accreted envelope. Here we examine the cooling evolution of neutron stars with polar field strengths ranging from 0 to 10^{16} G. The zero field calculation is based on the envelope model of

¹Presidential Faculty Fellow

Hernquist & Applegate (1984). The results of Hernquist (1985) describe the model envelope for 10^{13} G, and the ultramagnetized cases ($10^{14} - 10^{16}$ G) are calculated from the models of Heyl & Hernquist (1997a).

We choose a simple cooling model with the modified URCA process (Shapiro & Teukolsky 1983) and photon cooling competing (Heyl & Hernquist 1997b). Both the envelope calculations and the cooling calculations are performed in the frame of the surface; consequently, the observed effective temperature for a given core temperature is proportional to $g_s^{1/4}$ (g_s is the gravitation acceleration at the surface) and $(1 + z_g)^{-1}$ where

$$(1 + z_g)^{-1} = \sqrt{1 - \frac{2GM}{Rc^2}}. \quad (1)$$

To obtain the effective temperature as observed at infinity for a given equation of state (*i.e.* M and R), these two corrections must be performed. Additionally, since for a magnetized neutron star, the thermal emission is not isotropic along the surface, we present the flux-weighted mean $\bar{T}_{\text{eff}}$ over the entire surface.

Figure 1 depicts the thermal evolution for several values of the magnetic field at the pole. We have assumed that $g_s = 10^{14}$ cm/s². The horizontal line gives the value of $T_{\text{bb}} = 8 \times 10^5$ K found by Haberl et al. (1997). Depending on the size and mass of the neutron star, this line may move up or down slightly; however, this will not strongly affect the age estimate because temperature drops quickly with age for $t \gtrsim 3 \times 10^5$ yr. Additionally, we see that near this temperature, neutrino cooling and photon cooling both contribute; consequently, the temperature at this age does not depend strongly on field strength or composition.

From these evolutionary tracks, we find that the age of RX J0720.4-3125 may range from 1.2×10^5 yr for a 10^{14} G field with a hydrogen envelope to 3.6×10^5 yr for an unmagnetized neutron star. The intense fields yield ages intermediate to these. We will assume a polar field of 10^{14} G and an iron envelope yielding an age estimate of 3.3×10^5 yr. With this age, we can obtain an estimate of the magnetic field of the neutron star, if we assume that it has spun down by magnetic dipole radiation (Shapiro & Teukolsky 1983),

$$t = \frac{3Ic^3}{B_p^2 R^6 \sin^2 \alpha \Omega_0^2}. \quad (2)$$

Solving for the magnetic field strength yields,

$$B_p \sin \alpha = \sqrt{\frac{3Ic^3}{t}} \frac{P}{2\pi R^3} = 9.3 \times 10^{13} \text{G} \left(\frac{M}{M_\odot} \right)^{1/2} \left(\frac{R}{10^6 \text{cm}} \right)^{-2} \left(\frac{t}{3.3 \times 10^5 \text{yr}} \right)^{-1/2} \quad (3)$$

This age estimate also yields a value for $\dot{P}$. For spin down by magnetic dipole radiation,

$$T = \frac{P}{\dot{P}} = 2t \quad (4)$$

which yields $\dot{P} = 4.3 \times 10^{-13}$. This is consistent with the observed $\dot{P} = -6.0 - 0.8 \times 10^{-12}$.

3. Discussion

We present a simple model for RX J0720.4-3125. Its emission originates from neutron-star cooling through a magnetized envelope. We derive a magnetic field $\sim 10^{14}$ G, an age $\sim 3 \times 10^5$ yr and a period derivative of 4×10^{-13} . The primary contrast between the predictions of this model and those of Wang (1997) is the value of $\dot{P}$.

Wang (1997) argues that RX 0720.4-3125 is an old object, $t \gtrsim 10^9$ yr. It initially had a much stronger field ($\sim 10^{12}$ G) than at present. Before decaying to $\sim 10^{10}$ G, the magnetic field spun down the neutron star to its present rate. Wang argues that accretion powers the present emission and that a value of $\dot{P} \lesssim 10^{-16}$ is consistent with this field-decay model.

Our prediction of $\dot{P}$ lies just within the current limits which Haberl et al. (1997) obtained over a three-year baseline. An further round of observations within the next several years could determine whether RX J0720.4-3125 is a middle-aged, isolated cooling neutron star or an ancient neutron star accreting from the interstellar medium.

The work was supported in part by a National Science Foundation Graduate Research Fellowship, the NSF Presidential Faculty Fellows program and Cal Space grant CS-12-97.

REFERENCES

- Haberl, F., Motch, C., Buckley, D. A. H., Zickgraf, F.-J. & Pietsch, W. 1997, *A&A*, in press
- Hernquist, L. & Applegate, J. H. 1984, *ApJ*, **287**, 244.
- Hernquist, L. 1985, *MNRAS*, **213**, 313.
- Heyl, J. S. & Hernquist, L. 1997a, *Almost Analytic Models of Ultramagnetized Neutron Star Envelopes*, submitted to *MNRAS*
- Heyl, J. S. & Hernquist, L. 1997b, *The Thermal Evolution of Ultramagnetized Neutron Stars*, submitted to *ApJL*
- Heyl, J. S. & Hernquist, L. 1997c, *Powering Anomalous X-ray Pulsars by Neutron Star Cooling*, submitted to *ApJL*
- Mereghetti, S. & Stella, L. 1995, *ApJL*, **442**, 17.
- Shapiro, S. L. & Teukolsky, S. A. 1983, *Black Holes, White Dwarfs, and Neutron Stars*, Wiley-Interscience, New York
- Van Paradijs, J., Taam, R. E. & van den Heuvel, E. P. J. 1995, *A&A*, **299**, 41.
- Wang, J. C. L. 1997, *Evidence for magnetic field decay in RX J0720.4-3125*, astro-ph/9706288, submitted to *ApJL*




Fig. 1.— The curves trace the cooling evolution of isolated neutron stars by the modified URCA process and the radiation of photons from the surface for several field strengths with iron and hydrogen envelopes. The horizontal line traces the observed surface temperature of RX J0720.4-3125 of $8 \times 10^5 \text{ K}$.